

Pressure-driven Instabilities in Cylindrical Geometry: A New General Criterion

Pierre-Yves Longaretti*

Laboratoire d'Astrophysique de Grenoble, BP 53X, F-38041, Grenoble, France

(Dated: 03/20/03)

A new criterion for pressure-driven interchange instabilities in cylindrical geometry is derived, based on an alternate use of the Energy Principle. This criterion is inequivalent to Suydam's criterion and does not contain the magnetic shear. In fact, it is shown that Suydam's criterion relates to the instability of the slow magnetosonic branch, while the present criterion relates to the Alfvénic one, which is the most dangerous of the two. These findings imply that pressure-driven modes often exist even if Suydam's criterion is satisfied by a large margin.

PACS numbers: 47.65.+a; 52.30.Cv; 52.35.Py; 95.30.Qd

MHD instabilities are usually divided into pressure-driven and current-driven, depending on whether the destabilizing agent results from the interplay between the gas pressure gradient and the field-line curvature, or from the component of the current parallel to the magnetic field. The stability in cylindrical geometry with respect to both types of processes is an old and well-trodden subject. It was actively studied in the 60s and 70s, both in its own sake and as an idealization of more complex geometries for fusion devices. Important general results have been established relatively early on, and are now part of the standard lore on the subject [1]; among these Suydam's criterion [2] for interchange pressure-driven modes in an arbitrary screw pinch is directly related to the present work. Later developments have mostly been concerned with the stability properties of particular configurations such as tokamaks or reversed-field pinches.

In addition to fusion devices, astrophysical jets constitute another class of physical systems for which the existence of purely MHD unstable modes in cylindrical geometry is extremely relevant. However, there are only a handful of papers on current-driven instabilities in jets ([3, 4, 5], and references therein), and only a couple of papers on pressure-driven instabilities [6, 7]. This surprisingly small level of activity reflects the fact that the Kelvin-Helmholtz instability has long been thought to be the most important and dangerous for jet survival, a conjecture somewhat belied by recent numerical results [8].

Suydam's criterion stresses that the magnetic shear is an important stabilizing factor in cylindrical MHD columns. Indeed, although this criterion is only a necessary condition for stability, it is well-known that the magnetic shear plays an important role in the ability of nuclear fusion devices to survive the catastrophic disruption usually produced by MHD instabilities. However, it has also long been noticed that pressure-driven unstable modes are easily found in MHD equilibria, with growth-rates nearly independent of their wavenumber, independently of whether Suydam's criterion is satisfied or not (see, e.g., Ref. [9]).

One of the objectives of this Letter is to provide a theoretical explanation for this fact, and to show that the role of pressure-driven interchange modes has probably been underestimated in fusion physics. This is done through the derivation of a new general criterion for the stability of pressure-driven modes, and by a discussion of the physical meaning of this new criterion as well as Suydam's. The criterion is obtained from an alternate use of the MHD Energy Principle [1, 10, 11].

For static equilibria, the MHD perturbation equations can be written as

$$-\rho_o \omega^2 \xi = -\nabla \delta P_* + \delta \mathbf{T} \equiv \mathbf{F}(\xi), \quad (1)$$

where $\delta \mathbf{T}$ is the perturbation of the magnetic tension, and δP_* the total pressure perturbation, and ξ the fluid element displacement from equilibrium (a Fourier transform in time has been performed). Specializing to cylindrical equilibria, it is convenient to project this perturbation equation in the three directions $\mathbf{e}_r$, $\mathbf{e}_\parallel$, $\mathbf{e}_\perp$ (i.e., the radial direction, the direction parallel to the unperturbed magnetic field, and the remaining one required to define an orthonormal frame); furthermore the displacement can be chosen as a Fourier mode:

$$\xi = \xi(r) \exp[i(\omega t - m\theta - k_z z)]. \quad (2)$$

This choice reflects the fact that the equilibrium quantities (the magnetic field line helical structure, the magnetic field components and the gas pressure) depend only on radius. The equations for the parallel and perpendicular displacement $\xi_\parallel$ and $\xi_\perp$ can be solved exactly as a function of ω^2 and ξ_r [1, 12, 13]:

$$\xi_{\parallel,\perp} = \xi_{\parallel,\perp}(\omega^2, \xi_r), \quad (3)$$

leading to an exact second order equation for the radial displacement [1, 12, 13]:

$$-\rho \omega^2 \xi = \frac{d}{dr} \left[A \frac{d}{dr} (r \xi_r) \right] + C^* r \xi_r \equiv F_r(\xi_r). \quad (4)$$

In this expression, A and C^* depend in a complicated way on the equilibrium quantities and on the mode frequency ω .

Let us define

$$\delta W_r(\xi_r^*, \xi_r) = -\frac{1}{2} \int \xi_r^* F_r(\xi) d^3 r, \quad (5)$$

and

$$\delta K_r(\xi_r^*, \xi_r) = \frac{1}{2} \int \rho_o |\xi_r|^2 d^3 r. \quad (6)$$

where, for any choice of ξ_r , the remaining displacement components $\xi_{||}$ and $\xi_{\perp}$ are specified by Eqs. (3), and $\omega^2 = \omega^2[\xi_r]$ is chosen to be the formal solution of the implicit equation

$$\omega^2 = \frac{\delta W_r}{\delta K_r}. \quad (7)$$

It can be shown from the standard form of the Energy Principle that, if there exists a total displacement such that $\delta W_r < 0$, then the system is unstable; furthermore, ω^2 , as given by Eq. (7), also satisfies $\omega^2 = \delta W / \delta K$, where δW and δK are the usual potential and kinetic energy of the standard Energy Principle [13].

Obviously, this reexpression of the Energy Principle is useful only if Eq. (7) can be solved for ω^2 , which is in general a rather daunting task. Fortunately, an interesting and useful simplification of F_r is suggested by the ballooning asymptotic ordering. In particular, let us look for trial displacements where m, k_z satisfy $k_{\perp} L_o \gg 1$ while $k_{||} L_o \ll 1$ (L_o is the characteristic scale of equilibrium gradients). In other words, one considers displacements with large wavenumbers, except in the direction of the unperturbed magnetic field. Because of the magnetic shear, this imposes that the displacement must be nonvanishing only in the close vicinity to the magnetic resonance associated to m, k , and implicitly defined by $k_{||} = 0$. Furthermore, let us make the assumption that one can find a trial radial displacement ξ_r such that the frequency ω^2 which is the solution of Eq. (7) obeys the ballooning ordering, i.e. $\omega_f^2 \gg |\omega^2|$ ($\sim v_A^2/L_o^2, c_s^2/L_o^2 \gg v_A^2 k_{||}^2, c_s^2 k_{||}^2$, where $\omega_f^2 \simeq (v_A^2 + c_s^2) k_{\perp}^2$ is the fast magnetosonic frequency. This assumption is easily checked a posteriori.

With these approximations, $k_{||}$ can be expanded to first order with respect to the distance to the magnetic resonance r_o , to read

$$k_{||} = -k_{\perp} \frac{B_{\theta} B_z}{B_o^2} \left(\frac{r - r_o}{r} \right) \frac{d \ln |h|}{d \ln r}, \quad (8)$$

where $h \equiv r B_z / B_{\theta}$ is the pitch length (its logarithmic derivative is the magnetic shear), and B_{θ}, B_z, B_o are the

equilibrium azimuthal, vertical, and total magnetic field, respectively. The width of the region where the radial displacement is nonvanishing can be quantified from the constraint that $k_{||} L_o$ is small enough (see below) with the help of this expression. For large $k_{\perp} L_o$, this width is narrow enough so that one can neglect the variations with radius of all quantities (except $k_{||}$) in the corresponding region.

From Eq. (8) (and the approximations just described), the quantity C^* reduces to [13]

$$C^* = -\frac{2B_{\theta}^2}{r^3} \left(\frac{r P_o'}{B_o^2} \right) - \frac{4\beta B_{\theta}^4}{(1+\beta)r^3 B_o^2} \equiv \frac{\omega_c^2}{r}. \quad (9)$$

In this expression, the system of units is chosen such that $\mu_o = 1$, and the plasma β parameter is defined as $\beta \equiv c_s^2/v_A^2$, i.e. as the square of the ratio of the adiabatic sound speed (for an adiabatic equation of state) to the Alfvén speed; this definition differs from the standard one by a factor $\gamma/2$ (γ is the adiabatic index).

With these simplifications, Eq. (7) now reads

$$\omega^2 \int \rho_o |\xi_r|^2 = \int A r \left| \frac{d\xi_r}{dr} \right|^2 dr + \omega_c^2 \int \rho_o |\xi_r|^2 dr, \quad (10)$$

where, to leading order, A is given by

$$A = -\frac{\rho_o}{r} \frac{\omega^2}{k_{\perp}^2}. \quad (11)$$

Solving for ω^2 yields:

$$\omega^2 = \omega_c^2 \frac{\int_{r_o-\delta}^{r_o+\delta} |\xi_r|^2 dr}{\int_{r_o-\delta}^{r_o+\delta} (|\xi_r|^2 + 1/k_{\perp}^2 |d\xi_r/dr|^2) dr}, \quad (12)$$

where ω_c^2 is defined in Eq. (9). The quantity δ appearing in Eq. (12) measures the width of the region where the radial displacement is nonvanishing. It must be chosen such that the constraint

$$v_A^2 k_{||}^2, c_s^2 k_{||}^2 \lesssim |\omega^2| \quad (13)$$

is satisfied, with the help of Eq. (8) (if the displacement is non zero in regions where $v_A^2 k_{||}^2, c_s^2 k_{||}^2 \gtrsim |\omega^2|$, it produces positive contributions to the potential energy δW_r [13]). It is apparent that any displacement satisfying this constraint makes $\omega^2 < 0$ provided that

$$\omega_c^2 = v_A^2 \left(-2\beta \kappa_{\rho} \kappa_c + \frac{4\beta \kappa_c^2}{1+\beta} \right) < 0. \quad (14)$$

In Eq. (14), $\kappa_c \equiv \mathbf{e}_r \cdot [(\mathbf{e}_{||} \cdot \nabla) \mathbf{e}_{||}] = -B_{\theta}^2/(B_o^2 r)$ is an algebraic measure of the field line inverse radius of curvature,

while $\kappa_\rho = (d\rho_o/dr)/\rho_o$ is a measure of the pressure gradient scale (an adiabatic equation of state is assumed). This reexpression of ω_c^2 allows us to recognise the first term as the usual destabilizing term due to the gas pressure and the field line curvature, while the second (stabilizing) term arises from the plasma compression. As a matter of fact, this criterion generalizes a result due to Kadomtsev for the sausage ($m = 0$) mode in a Z-pinch [14].

The reader can easily convince himself that the condition required earlier on ω^2 is satisfied for reasonable choices of ξ_r if $\omega_c^2 \neq 0$, thereby insuring the self-consistency of the result. In the whole procedure, and in contrast to common usage, neither the incompressibility condition nor marginal stability has been assumed (actually, the two conditions are known to be tied to one another). It is precisely because of this feature that it was possible to derive the criterion. Note that the usual reasoning according to which the most dangerous modes are incompressible is valid only if incompressible modes with finite growth rates can be found, which is not the case for pressure-driven modes in MHD static equilibria, as just pointed out.

Within the ballooning asymptotic ordering, there is another well-known limiting case of direct interest to the purpose of this Letter, that make Eq. (4) analytically tractable: namely, $\omega_f^2 \gg v_A^2/L_o^2, c_s^2/L_o^2 \gg v_A^2 k_\parallel^2, c_s^2 k_\parallel^2 \gg |\omega^2| \approx 0$. This limit allows us to recover Suydam criterion. Indeed, in this limit, the fast variation of $k_\parallel$ with radius dominates over all others, and after some algebra, Eq. (4) reduces to

$$\frac{d}{dr} \left(\frac{r B_o^2 k_\parallel^2}{k^2} \frac{d\xi_r}{dr} \right) - \frac{2k_z^2 P_o'}{k^2} \xi_r = 0, \quad (15)$$

where $k \equiv (k_z^2 + m^2/r^2)^{1/2}$. With the help of Eq. (8), this equation gives the behavior of ξ as a function of radius as $\xi \propto [(r - r_o)/r_o]^p$; p is real when

$$r B_z^2 \left(\frac{h'}{h} \right)^2 + 8P_o' > 0, \quad (16)$$

p is complex otherwise. In this last case, it is possible to show, with the help of the Energy Principle, that the fluid is unstable (see, e.g., Ref. [1] for details). Therefore, Eq. (16) (Suydam's criterion [2]) constitutes a necessary condition for stability.

This stresses the fact that both the new instability criterion Eq. (14), and Suydam's necessary stability criterion Eq. (16) follow from Eq. (4), but in different limits, and are not mutually exclusive. A deeper insight into the relation between these two criteria is obtained by considering the low magnetic shear limit. In this case, a WKB approximation can be made on ξ_r with radial wavenumber k_r satisfying $|k_\perp r| \gg |k_r r| \gg 1$, so that the term

involving the derivative of ξ_r can be neglected in Eq. (4). Assuming $\omega_f^2 \gg |\omega^2|, v_A^2 k_\parallel^2, c_s^2 k_\parallel^2$, Eq. (4) reduces to the following dispersion relation [7, 13] [23]

$$\omega^4 - \left[k_\parallel^2 (v_A^2 + v_{SM}^2) + \omega_c^2 \right] \omega^2 + k_\parallel^2 (k_\parallel^2 - 2\beta\kappa_\rho\kappa_c) v_{SM}^2 v_A^2 = 0, \quad (17)$$

where $v_{SM}^2 \simeq \beta v_A^2 / (1 + \beta)$ is the slow magnetosonic square speed.

A necessary and sufficient condition of instability is

$$k_\parallel^2 < 2\beta\kappa_\rho\kappa_c, \quad (18)$$

which requires in particular $2\beta\kappa_\rho\kappa_c > 0$, a very-well known constraint for the existence of pressure-driven instabilities. Note that Eq. (18) generalizes a criterion derived by Kadomtsev for the $m = 1$ mode [6, 7, 14]. In the limit $k_\parallel \rightarrow 0$, there is a small and a large root to the dispersion relation Eq. (17), which, to leading order in $k_\parallel L_o$ are approximately given by

$$\omega_-^2 \simeq -v_{SM}^2 \frac{2\beta\kappa_\rho\kappa_c v_A^2}{\omega_c^2} k_\parallel^2, \quad (19)$$

and

$$\omega_+^2 \simeq \omega_c^2, \quad (20)$$

Furthermore, in the limit $\kappa_\rho, \kappa_c \rightarrow 0$, the large and small roots reduce to the standard Alfvén and slow magnetosonic root, respectively. Only one of these roots is unstable at a time, depending on the sign of ω_c^2 .

Note that the large root ω_+^2 satisfies the ordering assumed in the derivation of the new criterion; furthermore, for displacements ξ_r satisfying the WKB approximation assumed in the derivation of Eq. (17), Eq. (12) gives back Eq. (20). Reversely, the large root mode cannot satisfy the ordering needed to rederive Suydam's criterion, in contrast to small root modes.

These arguments show that Suydam's criterion relates to the stability of the slow magnetosonic branch, while the newly derived criterion relates to the stability of the Alfvénic one, a conclusion that can also be reached by an analysis of the global modes behavior [13]. Of the two branches, the Alfvénic one the most dangerous, because it has the fastest growth rate, so that the new criterion is probably the more relevant of the two to ascertain the stability of MHD cylindrical equilibria.

The practical usefulness of this new criterion also follows from the fact that $\omega^2 \lesssim \omega_c^2/2$ always provides a rather precise order of magnitude estimate of the unstable pressure modes growth rates, even for low wavenumbers (down to $m = 1$, in fact [15]); this feature also explains why the most unstable pressure-driven modes have

a growth rate nearly independent of their wavenumber. Quite surprisingly, the new criterion does not contain the magnetic shear. In fact, the magnetic shear only role is to limit the mode width through Eq. (13) [13]; it is because of this feature, and not because of its influence on the growth rate, that the magnetic shear can limit the disruptive power of Alfvénic pressure-driven instabilities.

Under which conditions can one expect Eq. (14) to be satisfied ?

It is well-known that, in a Z-pinch equilibrium, $\omega_c^2 > 0$ is a constraint which is difficult to realize in practice. In more realistic equilibria, however, one expects the longitudinal component of the equilibrium magnetic field to decrease with increasing radius, so that the gas pressure-gradient is relatively weaker in such a configuration than in a Z-pinch equilibrium, a feature which stabilizes the plasma, unless the pressure gradient occurs only on a rather small radial scale. In fact, this is precisely what happens in some of the most commonly considered tokamaks or RFP configurations. Furthermore, fusion devices are collisionless over the time-scales under which they usually operate; the simplest (albeit crude) way to account for this is to set the adiabatic index $\gamma = 0$ in the problem, which suppresses the compression stabilizing term. Therefore, either for one or the other reason, pressure-driven interchange modes are necessarily present in fusion devices' plasmas, down to low azimuthal wave-numbers $m \gtrsim 1$ (in any case, large wavenumber modes should be stabilized by finite Larmor radius effects); furthermore, the growth rates are comparable to the Alfvén time-scale once the plasma β is of the order of a significant fraction of unity. These features strongly suggest that the role of pressure-driven interchange modes in the generation of the sawtooth oscillations and edge localized modes in tokamaks has been underestimated, a possibility which will be more fully explored elsewhere. These modes might also play a role in triggering the edge turbulent transport in fusion devices; on the other hand, it is now well-known that a velocity shear can both stabilize pressure-driven modes [16] and reduce the MHD-driven edge transport [17]. A velocity shear can also produce turbulent transport by itself through hydrodynamical processes, but the flow curvature and the Coriolis force can severely limit the induced turbulent transport [18, 19].

In what concerns astrophysical jets, most models studied in the literature are cold, i.e., the equilibrium is not provided by the balance between the pressure gradient and the magnetic tension due to the azimuthal field, but by a balance between the inertial force due to rotation

and the magnetic tension [20, 21, 22]. Destabilization by the rotation inertial force is expected in this situation [7], in spite of the stabilizing action of the Coriolis force, with features very similar to the ones described above. A more extensive study of rotation-driven instabilities in astrophysical jets will be presented elsewhere, while a comparison of usual pressure-driven instabilities with the Kelvin-Helmholtz instability is performed in Ref. [15].

* Electronic address: pyl@obs.ujf-grenoble.fr;
URL: <http://www-laog.obs.ujf-grenoble.fr/~pyl/>

- [1] J.P. Freidberg, *Ideal Magnetohydrodynamics*, Plenum Press, New York (1987).
- [2] B.R. Suydam. In *Proc. of the Second U.N. Internat. Conf. on the Peaceful Uses of Atomic Energy*, United Nations, Geneva, 31, 157 (1958).
- [3] S. Appl, and M. Camenzind, *Astron. Astrophys.* 256, 354 (1992).
- [4] S. Appl, *Astron. Astrophys.*, 314, 995 (1996).
- [5] S. Appl, T. Lery, and H. Baty, *Astron. Astrophys.*, 355, 818 (2000).
- [6] M.C. Belgelman, *Astrophys. J.*, 493, 291 (1998).
- [7] E. Kersalé, P.-Y. Longaretti, and G. Pelletier, *Astron. Astrophys.*, 363, 1166 (2000).
- [8] D. Ryu, T.W. Jones, and A. Frank, *Astrophys. J.*, 545, 475 (2000).
- [9] D. Merlin, S. Ortolani, R. Cappagnella, and M. Scapin, *Nucl. Fusion*, 29, 1153 (1989).
- [10] I.B. Bernstein, E.A. Frieman, M.D. Kruskal, and R.M. Kulsrud, *Proc. R. Soc. London*, A 244, 17 (1958).
- [11] G. Laval, C. Mercier, and R. Pellat, *Nucl. Fusion*, 5, 156 (1965).
- [12] K. Hain, and R. Lüster, *Z. Naturforsch. Teil A*, 13, 936 (1958).
- [13] P.-Y. Longaretti. *In preparation*.
- [14] B.B. Kadomtsev. In *Reviews of Plasma Physics*, ed. M.A. Leontovitch, Consultants Bureau, New York, Vol. II (1966).
- [15] P.-Y. Longaretti, and H. Baty. *Submitted to Astron. Astrophys.*
- [16] T. Chiueh, *Phys. Rev. E*, 54, 5632 (1996).
- [17] P. W. Terry, *Rev. Modern Phys.*, 72, 109 (2000).
- [18] P.-Y. Longaretti, *Astrophys. J.*, 576, 587 (2002).
- [19] P.-Y. Longaretti. *Submitted to Phys. Rev. Lett.*
- [20] G. Pelletier, R.E. Pudritz, *Astrophys. J.*, 394, 117 (1992).
- [21] F. Casse, and J. Ferreira, *Astron. Astrophys.*, 353, 1115 (2000).
- [22] J. Ferreira, *Astron. Astrophys.*, 319, 340 (1997).
- [23] There is an algebraic mistake in Ref. [7], but the results of this paper are valid in the limit of negligible magnetic shear considered here.